INVESTIGATOR'S ANNUAL REPORT

National Park Service

All or some of the information provided may be available to the public

| Reporting Year: | Park: Shenandoah NP |
|--|--|
| | |
| Principal Investigator: Richard Tollo | Office Phone: (202)994-6960 |
| Renard Torio | |
| | Email: |
| | rtollo@gwu.edu |
| Address: | Office Fax: |
| George Washington University | (202)994-0450 |
| Department of Earth and | |
| Environmental Sciences Washington, DC 20052 US | |
| Additional investigators or key field assistants (first name, last name, office phone, office email): | |
| No co-investigators | |
| Permit#: | |
| SHEN1996ARWP | |
| Park-assigned Study Id. #: unknown | |
| Project Title: Geologic Evolution Of Mesoproterozioc Basement, Blue Ridge Province, Shenandoah National Park, Virginia (N-133B) | |
| Permit Start Date: | Permit Expiration Date |
| Jan 01, 1998 | Jan 01, 1998 |
| Study Start Date: Jan 01, 1996 | Study End Date Jan 01, 1996 |
| Study Status: Completed | |
| Activity Type: Research | |
| Subject/Discipline: Geochemistry (inc. Minerals / Petrology) | |
| Objectives: | |
| The primary objective of this study is the determination of detailed geologic and geochronologic relationships characterizing Mesoproterozoic metamorphic basement units exposed in that portion of the Blue Ridge province located in Shenandoah National Park, Virginia. The project involves detailed field mapping and an integrated program of petrologic, geochemical and isotopic analyses designed to elucidate petrologic, geochemical, and temporal aspects of the Grenville orogeny. | |
| Findings and Status: | |
| Work completed to date includes: (1) detailed field mapping in two portions of the study area, (2) sample collection and initial processing, and (3) | |
| | |
| Work completed to date includes: (1) detailed field mapping in two portions petrographic analysis of selected rock samples. Field mapping and petrograunits exposed between Mary's Rock and Old Rag Mountain. These units inc | phic investigation have confirmed the presence of at least three lithologic |
| petrographic analysis of selected rock samples. Field mapping and petrogra units exposed between Mary's Rock and Old Rag Mountain. These units inc pyroxene charnockite, and (3) leucocratic charnockite. The discovery of suc | phic investigation have confirmed the presence of at least three lithologic clude: (1) garnet-bearing chanockitic gneiss, (2) amphibole-bearing two-ch lithologically diverse units within a relatively small area underscores the |
| petrographic analysis of selected rock samples. Field mapping and petrograunits exposed between Mary's Rock and Old Rag Mountain. These units inc pyroxene charnockite, and (3) leucocratic charnockite. The discovery of suc geological complexity of the region and further indicates that much geologic orogeny has been heretofore overlooked. Geochemical analyses of bulk roc | phic investigation have confirmed the presence of at least three lithologic clude: (1) garnet-bearing chanockitic gneiss, (2) amphibole-bearing two-ch lithologically diverse units within a relatively small area underscores the evidence pertaining to the nature, style, and significance of the Grenville k samples and chemical analyses of mineral samples are planned for Spring |
| petrographic analysis of selected rock samples. Field mapping and petrograunits exposed between Mary's Rock and Old Rag Mountain. These units inc pyroxene charnockite, and (3) leucocratic charnockite. The discovery of suc geological complexity of the region and further indicates that much geologic orogeny has been heretofore overlooked. Geochemical analyses of bulk roc 1997 and are intended to produce further evidence bearing on the nature of C | phic investigation have confirmed the presence of at least three lithologic clude: (1) garnet-bearing chanockitic gneiss, (2) amphibole-bearing two-ch lithologically diverse units within a relatively small area underscores the evidence pertaining to the nature, style, and significance of the Grenville k samples and chemical analyses of mineral samples are planned for Spring Grenville-age magmas and the temperature-pressure conditions at which |
| petrographic analysis of selected rock samples. Field mapping and petrograunits exposed between Mary's Rock and Old Rag Mountain. These units inc pyroxene charnockite, and (3) leucocratic charnockite. The discovery of suc geological complexity of the region and further indicates that much geologic orogeny has been heretofore overlooked. Geochemical analyses of bulk roc | phic investigation have confirmed the presence of at least three lithologic clude: (1) garnet-bearing chanockitic gneiss, (2) amphibole-bearing two-ch lithologically diverse units within a relatively small area underscores the evidence pertaining to the nature, style, and significance of the Grenville k samples and chemical analyses of mineral samples are planned for Spring Grenville-age magmas and the temperature-pressure conditions at which |
| petrographic analysis of selected rock samples. Field mapping and petrograunits exposed between Mary's Rock and Old Rag Mountain. These units inc pyroxene charnockite, and (3) leucocratic charnockite. The discovery of suc geological complexity of the region and further indicates that much geologic orogeny has been heretofore overlooked. Geochemical analyses of bulk roc 1997 and are intended to produce further evidence bearing on the nature of C basement formation occurred. Isotopic analyses of zircons and monazites fr | phic investigation have confirmed the presence of at least three lithologic clude: (1) garnet-bearing chanockitic gneiss, (2) amphibole-bearing two-th lithologically diverse units within a relatively small area underscores the evidence pertaining to the nature, style, and significance of the Grenville k samples and chemical analyses of mineral samples are planned for Spring Grenville-age magmas and the temperature-pressure conditions at which om selected samples will be undertaken during Summer 1997 and will be |
| petrographic analysis of selected rock samples. Field mapping and petrogra units exposed between Mary's Rock and Old Rag Mountain. These units inc pyroxene charnockite, and (3) leucocratic charnockite. The discovery of suc geological complexity of the region and further indicates that much geologic orogeny has been heretofore overlooked. Geochemical analyses of bulk roc 1997 and are intended to produce further evidence bearing on the nature of C basement formation occurred. Isotopic analyses of zircons and monazites frused to determine ages of Grenville-related plutonism and metamorphism. For this study, were one or more specimens collected and removed from | phic investigation have confirmed the presence of at least three lithologic clude: (1) garnet-bearing chanockitic gneiss, (2) amphibole-bearing two-th lithologically diverse units within a relatively small area underscores the evidence pertaining to the nature, style, and significance of the Grenville k samples and chemical analyses of mineral samples are planned for Spring Grenville-age magmas and the temperature-pressure conditions at which om selected samples will be undertaken during Summer 1997 and will be |

| Fill out the following ONLY IF the National Park Service supported this project in this reporting year by providing money to a university or college | |
|--|--|
| Full name of college or university: | Annual funding provided by NPS to university or college this reporting year: |
| n/a | 0 |